



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEATHS DURING WEEK ENDED MARCH 1, 1919, IN CITIES.

The following table shows the registered deaths from all causes, and from pneumonia (all forms) and influenza combined, in certain large cities of the United States during the week ended March 1, 1919. The annual death rates per 1,000 population for the week and for the corresponding week of previous years are also shown.

The data are taken from the Weekly Health Index, March 4, 1919, issued by the Bureau of the Census, Department of Commerce. The populations used in computing the rates are estimated by the Bureau of the Census as of July 1, 1918.

Registered deaths and annual death rates per 1,000 population in certain large cities of the United States, week ended Mar. 1, 1919—Deaths from all causes, and from pneumonia (all forms) and influenza combined.

City.	Popula- tion, July 1, 1918, estimated.	Total deaths, all causes.	Annual death rate per 1,000.	Annual death rate for preceding years. ¹	Influenza and pneu- monia (all forms).	
					Number of deaths.	Annual rate per 1,000.
Albany, N. Y.	112,565	44	20.4	C 21.8	12	5.6
Atlanta, Ga.	201,732	72	18.6	C 19.1	—	—
Baltimore, Md.	² 669,981	272	21.2	A 20.2	66	5.1
Birmingham, Ala.	197,670	74	19.5	A 18.7	—	—
Boston, Mass.	785,245	279	18.5	A 19.1	70	4.6
Buffalo, N. Y.	473,229	195	21.5	C 17.2	44	4.9
Cambridge, Mass.	111,432	27	12.6	A 16.1	2	.9
Chicago, Ill.	2,596,681	867	17.4	A 17.4	230	4.6
Cincinnati, Ohio.	418,022	214	26.7	C 19.2	90	11.2
Cleveland, Ohio.	810,306	259	16.7	C 13.5	94	6.0
Columbus, Ohio.	225,296	86	19.9	C 13.9	27	6.2
Dayton, Ohio.	130,655	52	20.8	C 14.8	16	6.4
Denver, Colo.	—	96	—	—	—	—
Fall River, Mass.	128,392	31	12.6	C 15.8	13	5.3
Grand Rapids, Mich.	135,450	35	13.5	C 15.0	—	—
Indianapolis, Ind.	289,577	114	20.6	C 18.4	—	—
Kansas City, Mo.	313,785	165	27.4	C 18.9	55	9.1
Los Angeles, Calif.	568,495	147	13.5	A 14.6	14	1.3
Louisville, Ky.	242,707	116	24.9	C 25.1	34	7.3
Lowell, Mass.	109,081	52	24.9	A 20.7	13	6.2
Memphis, Tenn.	154,759	74	24.9	C 27.6	15	5.1
Milwaukee, Wis.	453,481	94	10.8	A 14.4	—	—
Minneapolis, Minn.	383,442	107	14.6	C 12.1	—	—
Nashville, Tenn.	119,215	61	26.7	C 19.7	19	8.3
Newark, N. J.	428,684	163	19.8	C 19.9	54	6.6
New Haven, Conn.	154,865	49	16.5	C 21.9	—	—
New Orleans, La.	382,273	163	22.2	A 22.1	30	4.1
New York, N. Y.	5,215,879	2,158	21.6	C 17.9	904	9.0
Oakland, Calif.	214,206	47	11.4	A 11.7	—	—
Omaha, Nebr.	180,264	40	11.6	C 12.4	—	—
Philadelphia, Pa.	1,761,371	682	20.2	³ 18.4	207	6.1
Pittsburgh, Pa.	593,303	287	25.2	C 19.5	134	11.8
Portland, Oreg.	—	70	—	—	7	—
Providence, R. I.	263,613	93	18.4	C 17.0	21	4.2
Richmond, Va.	160,719	60	19.5	C 23.4	10	3.2
Rochester, N. Y.	264,855	79	15.6	C 16.5	7	1.4
St. Louis, Mo.	779,951	257	17.2	C 17.6	81	5.4
St. Paul, Minn.	257,699	52	10.5	C 9.3	—	—
San Francisco, Calif.	478,530	182	19.8	C 18.4	21	2.3
Seattle, Wash.	—	71	—	—	12	—
Spokane, Wash.	—	19	—	—	—	—
Syracuse, N. Y.	161,404	67	21.6	C 16.2	—	—
Toledo, Ohio.	262,234	69	13.7	A 15.1	14	2.8
Washington, D. C.	401,681	163	21.2	A 20.5	35	4.5
Worcester, Mass.	173,650	60	18.0	C 15.9	8	2.4

¹ "A" indicates that the rate given is the average annual death rate per 1,000 population for the corresponding week of the years 1913 to 1917, inclusive. "C" indicates that the rate is the annual death rate per 1,000 population for the corresponding week of 1918.

² Population estimated as of July 1, 1919.

³ Rate is based on statistics of 1915, 1916, and 1917.